

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

Product datasheet for TA336445

Presenilin 2 (PSEN2) Mouse Monoclonal Antibody [Clone ID: 198C679.2.]

Product data:

Product Type:	Primary Antibodies
Clone Name:	198C679.2.
Applications:	WB
Recommend Dilution:	WB: 1:250-1:500
Reactivity:	Human
Host:	Mouse
lsotype:	IgM
Clonality:	Monoclonal
Immunogen:	This antibody was generated by immunizing mice with a synthetic peptide of human Presenilin-2 (PS-2).
Formulation:	Preservative: 0.1% Sodium Azide. Store at 4C short term. Aliquot and store at -20C long term. Avoid freeze-thaw cycles.
Concentration:	This product is unpurified. The exact concentration of antibody is not quantifiable.
Purification:	Tissue culture supernatant
Gene Name:	presenilin 2
Database Link:	<u>NP_036618 Entrez Gene 5664 Human</u>
Background:	Alzheimer's disease (AD) patients with an inherited form of the disease carry mutations in the presenilin proteins (PSEN1; PSEN2) or the amyloid precursor protein (APP). These disease- linked mutations result in increased production of the longer form of amyloid-beta (main
	component of amyloid deposits found in AD brains). Presenilins are postulated to regulate APP processing through their effects on gamma-secretase, an enzyme that cleaves APP. Also, it is thought that the presenilins are involved in the cleavage of the Notch receptor, such that they either directly regulate gamma-secretase activity or themselves are protease enzymes. Two alternative transcripts of PSEN2 have been identified.
Synonyms:	APP processing through their effects on gamma-secretase, an enzyme that cleaves APP. Also, it is thought that the presenilins are involved in the cleavage of the Notch receptor, such that they either directly regulate gamma-secretase activity or themselves are protease enzymes.
Synonyms: Protein Families:	APP processing through their effects on gamma-secretase, an enzyme that cleaves APP. Also, it is thought that the presenilins are involved in the cleavage of the Notch receptor, such that they either directly regulate gamma-secretase activity or themselves are protease enzymes. Two alternative transcripts of PSEN2 have been identified.



This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2020 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US

Product images:

MW (kDa) 200	
116 97	=
66	- 25
55	-
36	- 27
31	
21	- 12
14	- 22
6	- 20

Western Blot: Presenilin 2 Antibody (198C679.2.) TA336445 - analysis for Presenilin-2 using this antibody. A protein band of approximate molecular weight of 55 kDa was detected.

This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2020 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US