

Product datasheet for TA347061

TBL1 (TBL1X) Mouse Monoclonal Antibody [Clone ID: 4H2-D5-E9]

Product data:

Product Type:	Primary Antibodies
Clone Name:	4H2-D5-E9
Applications:	IF, WB
Recommend Dilution:	WB: 1:1000, IF: 1:100
Reactivity:	Human
Host:	Mouse
Isotype:	IgG2b
Clonality:	Monoclonal
Immunogen:	The immunogen for TBL1X antibody: purified recombinant human TBL1 protein fragments expressed in E.coli.
Formulation:	Purified mouse monoclonal antibody in PBS(pH 7.4) containing with 0.02% sodium azide and 50% glycerol.
Purification:	Affinity purified
Predicted Protein Size:	58 kDa
Gene Name:	transducin (beta)-like 1X-linked
Database Link:	NP_005638 Entrez Gene 6907 Human
Background:	The protein encoded by this gene has sequence similarity with members of the WD40 repeat-containing protein family. The WD40 group is a large family of proteins, which appear to have a regulatory function. It is believed that the WD40 repeats mediate protein-protein interactions and members of the family are involved in signal transduction, RNA processing, gene regulation, vesicular trafficking, cytoskeletal assembly and may play a role in the control of cytotypic differentiation. This encoded protein is found as a subunit in corepressor SMRT (silencing mediator for retinoid and thyroid receptors) complex along with histone deacetylase 3 protein. This gene is located adjacent to the ocular albinism gene and it is thought to be involved in the pathogenesis of the ocular albinism with late-onset sensorineural deafness phenotype. Four transcript variants encoding two different isoforms have been found for this gene. This gene is highly similar to the Y chromosome TBL1Y gene. [provided by RefSeq, Nov 2008]
Synonyms:	EBI; SMAP55; TBL1




[View online »](#)


Protein Families: Transcription Factors

Protein Pathways: Wnt signaling pathway

Product images:



Western blot detection of TBL1X in SW480 cell lysates using TBL1X mouse mAb (1:1000 diluted). Predicted band size: 58KDa. Observed band size: 58KDa.



Immunocytochemistry staining of HeLa cells fixed with 4% Paraformaldehyde and using anti-TBL1X mouse mAb (dilution 1:100).