

## Product datasheet for **TA500028**

### **Noggin (NOG) Mouse Monoclonal Antibody [Clone ID: OTI4C12]**

#### **Product data:**

Product Type:	Primary Antibodies
Clone Name:	OTI4C12
Applications:	IF, IHC, WB
Recommend Dilution:	WB 1:1000~2000, IHC 1:50, IF 1:100
Reactivity:	Human, Dog
Host:	Mouse
Isotype:	IgG2b
Clonality:	Monoclonal
Immunogen:	Human recombinant protein fragment corresponding to amino acids 28-232 of human noggin (NP_005441) produced in E.coli.
Formulation:	PBS (pH 7.3) containing 1% BSA, 50% glycerol and 0.02% sodium azide.
Concentration:	1 mg/ml
Purification:	Purified from mouse ascites fluids or tissue culture supernatant by affinity chromatography (protein A/G)
Predicted Protein Size:	23.7 kDa
Gene Name:	noggin
Database Link:	<a href="#">NP_005441</a> <a href="#">Entrez Gene 100683515 Dog</a> <a href="#">Entrez Gene 9241 Human</a>



[View online »](#)

**Background:**

Noggin is a secreted polypeptide which binds and inactivates members of the transforming growth factor-beta (TGF-beta) superfamily signaling proteins, such as bone morphogenetic protein-4 (BMP4). By diffusing through extracellular matrices more efficiently than members of the TGF-beta superfamily, it may have a principal role in creating morphogenic gradients. Noggin appears to have pleiotropic effect, both early in development as well as in later stages. The results of the mouse knockout of the ortholog suggest that noggin is involved in numerous developmental processes, such as neural tube fusion and joint formation. Recently, several dominant human NOG mutations in unrelated families with proximal symphalangism (SYM1) and multiple synostoses syndrome (SYNS1) were identified; both SYM1 and SYNS1 have multiple joint fusion as their principal feature, and map to the same region (17q22) as this gene. All of these mutations altered evolutionarily conserved amino acid residues. The amino acid sequence of this human gene is highly homologous to that of *Xenopus*, rat and mouse.

**Synonyms:**


SYM1; SYNS1; SYNS1A

**Protein Families:**




Druggable Genome, Secreted Protein


**Protein Pathways:**

TGF-beta signaling pathway


**Product images:**

HEK293T cells were transfected with the pCMV6-ENTRY control (Left lane) or pCMV6-ENTRY Nog ([RC205020], Right lane) cDNA for 48 hrs and lysed. Equivalent amounts of cell lysates (5 ug per lane) were separated by SDS-PAGE and immunoblotted with anti-Nog. Positive lysates [LY401672] (100ug) and [LC401672] (20ug) can be purchased separately from OriGene.





Anti-Nog mouse monoclonal antibody (TA500028) immunofluorescent staining of COS7 cells transiently transfected by pCMV6-ENTRY Nog ([RC205020]).



Immunofluorescent staining of A549 cells using anti-Nog mouse monoclonal antibody (TA500028).