

Product datasheet for TA500496


STK39 Mouse Monoclonal Antibody [Clone ID: OTI4H3]

Product data:


Product Type:	Primary Antibodies
Clone Name:	OTI4H3
Applications:	FC, IF, IHC, WB
Recommend Dilution:	WB 1:1000~2000, IHC 1:50, IF 1:50, FLOW 1:100
Reactivity:	Human
Host:	Mouse
Isotype:	IgG2b
Clonality:	Monoclonal
Immunogen:	Full length human recombinant protein of human STK39 (NP_037365) produced in HEK293T cell.
Formulation:	PBS (pH 7.3) containing 1% BSA, 50% glycerol and 0.02% sodium azide.
Concentration:	1 mg/ml
Purification:	Purified from mouse ascites fluids or tissue culture supernatant by affinity chromatography (protein A/G)
Predicted Protein Size:	59.3 kDa
Gene Name:	serine/threonine kinase 39
Database Link:	NP_037365 Entrez Gene 27347 Human
Background:	This gene encodes a serine/threonine kinase that is thought to function in the cellular stress response pathway. The kinase is activated in response to hypotonic stress, leading to phosphorylation of several cation-chloride-coupled cotransporters. The catalytically active kinase specifically activates the p38 MAP kinase pathway, and its interaction with p38 decreases upon cellular stress, suggesting that this kinase may serve as an intermediate in the response to cellular stress. [provided by RefSeq]
Synonyms:	DCHT; PASK; SPAK
Protein Families:	Druggable Genome, Protein Kinase




[View online »](#)

Product images:



HEK293T cells were transfected with the pCMV6-ENTRY control (Left lane) or pCMV6-ENTRY STK39 ([RC223981], Right lane) cDNA for 48 hrs and lysed. Equivalent amounts of cell lysates (5 ug per lane) were separated by SDS-PAGE and immunoblotted with anti-STK39. Positive lysates [LY402227] (100ug) and [LC402227] (20ug) can be purchased separately from OriGene.




Western blot analysis of extracts (35ug) from 9 different cell lines by using anti-STK39 monoclonal antibody (HepG2: human; HeLa: human; SVT2: mouse; A549: human; COS7: monkey; Jurkat: human; MDCK: canine; PC12: rat; MCF7: human).



Immunohistochemical staining of paraffin-embedded Carcinoma of Human thyroid tissue using anti-STK39 mouse monoclonal antibody. (Heat-induced epitope retrieval by 10mM citric buffer, pH6.0, 100°C for 10min, TA500496)



Immunofluorescent staining of COS7 cells using anti-STK39 mouse monoclonal antibody (TA500496).



HEK293T cells transfected with either [RC223981] overexpress plasmid (Red) or empty vector control plasmid (Blue) were immunostained by anti-STK39 antibody (TA500496), and then analyzed by flow cytometry.