

## Product datasheet for **TA500741**


### **AKR1A1 Mouse Monoclonal Antibody [Clone ID: OTI8E12]**

#### **Product data:**


<b>Product Type:</b>	Primary Antibodies
<b>Clone Name:</b>	OTI8E12
<b>Applications:</b>	FC, IHC, WB
<b>Recommend Dilution:</b>	WB 1:2000, IHC 1:50, Flow 1:100
<b>Reactivity:</b>	Human
<b>Host:</b>	Mouse
<b>Isotype:</b>	IgG1
<b>Clonality:</b>	Monoclonal
<b>Immunogen:</b>	Full length human recombinant protein of human AKR1A1 (NP_006057) produced in HEK293T cell.
<b>Formulation:</b>	PBS (pH 7.3) containing 1% BSA, 50% glycerol and 0.02% sodium azide.
<b>Concentration:</b>	0.58 mg/ml
<b>Purification:</b>	Purified from mouse ascites fluids or tissue culture supernatant by affinity chromatography (protein A/G)
<b>Predicted Protein Size:</b>	36.6 kDa
<b>Gene Name:</b>	aldo-keto reductase family 1 member A1
<b>Database Link:</b>	<a href="#">NP_006057 Entrez Gene 10327 Human</a>
<b>Background:</b>	This gene encodes a member of the aldo/keto reductase superfamily, which consists of more than 40 known enzymes and proteins. This member, also known as aldehyde reductase, is involved in the reduction of biogenic and xenobiotic aldehydes and is present in virtually every tissue. Alternative splicing of this gene results in two transcript variants encoding the same protein.
<b>Synonyms:</b>	ALDR1; ALR; ARM; DD3; HEL-S-6
<b>Protein Families:</b>	Druggable Genome
<b>Protein Pathways:</b>	Glycerolipid metabolism, Glycolysis / Gluconeogenesis, Metabolic pathways




[View online »](#)

**Product images:**


HEK293T cells were transfected with the pCMV6-ENTRY control (Left lane) or pCMV6-ENTRY AKR1A1 ([RC200302], Right lane) cDNA for 48 hrs and lysed. Equivalent amounts of cell lysates (5 ug per lane) were separated by SDS-PAGE and immunoblotted with anti-AKR1A1. Positive lysates [LY401826] (100ug) and [LC401826] (20ug) can be purchased separately from OriGene.




Immunohistochemical staining of paraffin-embedded Kidney tissue within the normal limits using anti-AKR1A1 mouse monoclonal antibody. (Heat-induced epitope retrieval by 10mM citric buffer, pH6.0, 100°C for 10min, TA500741, Dilution 1:50)



Immunohistochemical staining of paraffin-embedded liver tissue within the normal limits using anti-AKR1A1 mouse monoclonal antibody. (Heat-induced epitope retrieval by 10mM citric buffer, pH6.0, 100°C for 10min, TA500741, Dilution 1:50)



Immunohistochemical staining of paraffin-embedded Carcinoma of liver tissue using anti-AKR1A1 mouse monoclonal antibody. (Heat-induced epitope retrieval by 10mM citric buffer, pH6.0, 100°C for 10min, TA500741, Dilution 1:50)



Flow cytometric analysis of HeLa cells, using anti-AKR1A1 antibody (TA500741), (Red) compared to a nonspecific negative control antibody (TA50011) (Blue).