

## OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

## **Product datasheet for TA500843**

## Claudin 5 (CLDN5) Mouse Monoclonal Antibody [Clone ID: OTI1G4]

**Product data:** 

**Product Type:** Primary Antibodies

Clone Name: OTI1G4
Applications: FC, IF

**Recommend Dilution:** IF 1:100, FLOW 1:100

Reactivity: Human
Host: Mouse
Isotype: IgG1

Clonality: Monoclonal

Immunogen: Full length human recombinant protein of human CLDN5(NP\_003268) produced in HEK293T

cell

**Formulation:** PBS (pH 7.3) containing 1% BSA, 50% glycerol and 0.02% sodium azide.

Concentration: 1 mg/ml

**Purification:** Purified from mouse ascites fluids or tissue culture supernatant by affinity chromatography

(protein A/G)

Predicted Protein Size: 23 kDa
Gene Name: claudin 5

Database Link: NP 003268 Entrez Gene 7122 Human

**Background:** This gene encodes a member of the claudin family. Claudins are integral membrane proteins

and components of tight junction strands. Tight junction strands serve as a physical barrier to prevent solutes and water from passing freely through the paracellular space between

epithelial or endothelial cell sheets. Mutations in this gene have been found in patients with velocardiofacial syndrome. Alternatively spliced transcript variants encoding the same

protein have been found for this gene. [provided by RefSeq]

Synonyms: AWAL; BEC1; CPETRL1; TMVCF


**Protein Families:** Transmembrane

**Protein Pathways:** Cell adhesion molecules (CAMs), Leukocyte transendothelial migration, Tight junction






## **Product images:**



Anti-CLDN5 mouse monoclonal antibody (TA500843) immunofluorescent staining of COS7 cells transiently transfected by pCMV6-ENTRY CLDN5 ([RC207122]).



HEK293T cells transfected with either [RC207122] overexpress plasmid (Red) or empty vector control plasmid (Blue) were immunostained by anti-CLDN5 antibody (TA500843), and then analyzed by flow cytometry.