

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

Product datasheet for TA501284

ketohexokinase (KHK) Mouse Monoclonal Antibody [Clone ID: OTI4E9]

Product data:

Product Type: Primary Antibodies

Clone Name: OTI4E9

Applications: FC, IF, IHC, WB

Recommend Dilution: WB 1:2000, IHC 1:50, IF 1:100, FLOW 1:100

Reactivity: Human
Host: Mouse
Isotype: IgG1

Clonality: Monoclonal

Immunogen: Full length human recombinant protein of human KHK (NP_000212) produced in HEK293T

cell.

Formulation: PBS (pH 7.3) containing 1% BSA, 50% glycerol and 0.02% sodium azide.

Concentration: 0.66 mg/ml

Purification: Purified from mouse ascites fluids or tissue culture supernatant by affinity chromatography

(protein A/G)

Predicted Protein Size: 32.5 kDa

Gene Name: ketohexokinase

Database Link: NP 000212 Entrez Gene 3795 Human

Background: This gene encodes ketohexokinase that catalyzes conversion of fructose to fructose-1-

phosphate. The product of this gene is the first enzyme with a specialized pathway that catabolizes dietary fructose. Alternatively spliced transcript variants encoding different

isoforms have been identified. [provided by RefSeq]

Synonyms: ketohexokinase; ketohexokinase (fructokinase)


Protein Families: Druggable Genome

Protein Pathways: Fructose and mannose metabolism, Metabolic pathways







Product images:



HEK293T cells were transfected with the pCMV6-ENTRY control (Left lane) or pCMV6-ENTRY KHK ([RC202424], Right lane) cDNA for 48 hrs and lysed. Equivalent amounts of cell lysates (5 ug per lane) were separated by SDS-PAGE and immunoblotted with anti-KHK. Positive lysates [LY400082] (100ug) and [LC400082] (20ug) can be purchased separately from OriGene.




Immunohistochemical staining of paraffinembedded Human Kidney tissue within the normal limits using anti-KHK mouse monoclonal antibody. (Heat-induced epitope retrieval by 10mM citric buffer, pH6.0, 100°C for 10min, TA501284, Dilution 1:50)




Immunohistochemical staining of paraffinembedded Human liver tissue within the normal limits using anti-KHK mouse monoclonal antibody. (Heat-induced epitope retrieval by 10mM citric buffer, pH6.0, 100°C for 10min, TA501284, Dilution 1:50)





Anti-KHK mouse monoclonal antibody (TA501284) immunofluorescent staining of COS7 cells transiently transfected by pCMV6-ENTRY KHK ([RC202424]).



HEK293T cells transfected with either [RC202424] overexpress plasmid (Red) or empty vector control plasmid (Blue) were immunostained by anti-KHK antibody (TA501284), and then analyzed by flow cytometry.