

Product datasheet for **TA501529**

BTN1A1 Mouse Monoclonal Antibody [Clone ID: OTI7B5]


Product data:

Product Type:	Primary Antibodies
Clone Name:	OTI7B5
Applications:	FC, IF, IHC, WB
Recommend Dilution:	WB 1:200~500, IHC 1:150, IF 1:100, FLOW 1:100
Reactivity:	Human, Dog
Host:	Mouse
Isotype:	IgG2a
Clonality:	Monoclonal
Immunogen:	Full length human recombinant protein of human BTN1A1 (NP_001723) produced in HEK293T cell.
Formulation:	PBS (PH 7.3) containing 1% BSA, 50% glycerol and 0.02% sodium azide.
Concentration:	0.58 mg/ml
Purification:	Purified from mouse ascites fluids or tissue culture supernatant by affinity chromatography (protein A/G)
Predicted Protein Size:	58.8 kDa
Gene Name:	butyrophilin subfamily 1 member A1
Database Link:	NP_001723 Entrez Gene 100855656 Dog Entrez Gene 696 Human
Background:	Butyrophilin is the major protein associated with fat droplets in the milk. It is a member of the immunoglobulin superfamily. It may have a cell surface receptor function. The human butyrophilin gene is localized in the major histocompatibility complex (MHC) class I region of 6p and may have arisen relatively recently in evolution by the shuffling of exons between 2 ancestral gene families [provided by RefSeq].
Synonyms:	BT; BTN; BTN1
Protein Families:	Druggable Genome, Transmembrane




[View online »](#)


Product images:




HEK293T cells were transfected with the pCMV6-ENTRY control (Left lane) or pCMV6-ENTRY BTN1A1 ([RC223852], Right lane) cDNA for 48 hrs and lysed. Equivalent amounts of cell lysates (5 ug per lane) were separated by SDS-PAGE and immunoblotted with anti-BTN1A1. Positive lysates [LY400653] (100ug) and [LC400653] (20ug) can be purchased separately from OriGene.




Western blot analysis of extracts (35ug) from 9 different cell lines by using anti-BTN1A1 monoclonal antibody.




Immunohistochemical staining of paraffin-embedded Human colon tissue within the normal limits using anti-BTN1A1 mouse monoclonal antibody. (Heat-induced epitope retrieval by 10mM citric buffer, pH6.0, 100°C for 10min, TA501529)




Immunohistochemical staining of paraffin-embedded Carcinoma of Human kidney tissue using anti-BTN1A1 mouse monoclonal antibody. (Heat-induced epitope retrieval by 10mM citric buffer, pH6.0, 100°C for 10min, TA501529)




Immunohistochemical staining of paraffin-embedded Carcinoma of Human lung tissue using anti-BTN1A1 mouse monoclonal antibody. (Heat-induced epitope retrieval by 10mM citric buffer, pH6.0, 100°C for 10min, TA501529)




Immunohistochemical staining of paraffin-embedded Adenocarcinoma of Human ovary tissue using anti-BTN1A1 mouse monoclonal antibody. (Heat-induced epitope retrieval by 10mM citric buffer, pH6.0, 100°C for 10min, TA501529)




Immunohistochemical staining of paraffin-embedded Human pancreas tissue within the normal limits using anti-BTN1A1 mouse monoclonal antibody. (Heat-induced epitope retrieval by 10mM citric buffer, pH6.0, 100°C for 10min, TA501529)




Immunohistochemical staining of paraffin-embedded Human thyroid tissue within the normal limits using anti-BTN1A1 mouse monoclonal antibody. (Heat-induced epitope retrieval by 10mM citric buffer, pH6.0, 100°C for 10min, TA501529)




Immunohistochemical staining of paraffin-embedded Adenocarcinoma of Human endometrium tissue using anti-BTN1A1 mouse monoclonal antibody. (Heat-induced epitope retrieval by 10mM citric buffer, pH6.0, 100°C for 10min, TA501529)




Immunohistochemical staining of paraffin-embedded Human prostate tissue within the normal limits using anti-BTN1A1 mouse monoclonal antibody. (Heat-induced epitope retrieval by 10mM citric buffer, pH6.0, 100°C for 10min, TA501529)




Immunohistochemical staining of paraffin-embedded Carcinoma of Human bladder tissue using anti-BTN1A1 mouse monoclonal antibody. (Heat-induced epitope retrieval by 10mM citric buffer, pH6.0, 100°C for 10min, TA501529)




Anti-BTN1A1 mouse monoclonal antibody (TA501529) immunofluorescent staining of COS7 cells transiently transfected by pCMV6-ENTRY BTN1A1 ([RC223852]).



HEK293T cells transfected with either [RC223852] overexpress plasmid (Red) or empty vector control plasmid (Blue) were immunostained by anti-BTN1A1 antibody (TA501529), and then analyzed by flow cytometry.



Flow cytometric Analysis of HeLa cells, using anti-BTN1A1 antibody (TA501529), (Red), compared to a nonspecific negative control antibody ([TA50011]), (Blue).



Flow cytometric Analysis of Jurkat cells, using anti-BTN1A1 antibody (TA501529), (Red), compared to a nonspecific negative control antibody ([TA50011]), (Blue).