

## Product datasheet for TA501723


### SDR O (SDR9C7) Mouse Monoclonal Antibody [Clone ID: OTI4B5]

#### Product data:


Product Type:	Primary Antibodies
Clone Name:	OTI4B5
Applications:	FC, IHC, WB
Recommend Dilution:	WB 1:500, IHC 1:150, FLOW 1:100
Reactivity:	Human
Host:	Mouse
Isotype:	IgG1
Clonality:	Monoclonal
Immunogen:	Full length human recombinant protein of human SDR9C7 (NP_683695) produced in E.coli.
Formulation:	PBS (PH 7.3) containing 1% BSA, 50% glycerol and 0.02% sodium azide.
Concentration:	0.67 mg/ml
Purification:	Purified from mouse ascites fluids or tissue culture supernatant by affinity chromatography (protein A/G)
Predicted Protein Size:	35.1 kDa
Gene Name:	short chain dehydrogenase/reductase family 9C member 7
Database Link:	<a href="#">NP_683695 Entrez Gene 121214 Human</a>
Background:	This gene encodes a protein with similarity to the short-chain dehydrogenase/reductase (SDR) family but has not been shown to have retinoid or dehydrogenase activities. [provided by RefSeq]
Synonyms:	RDHS; SDR-O; SDRO
Protein Families:	Druggable Genome




[View online »](#)

**Product images:**


HEK293T cells were transfected with the pCMV6-ENTRY control (Left lane) or pCMV6-ENTRY SDR9C7 ([RC210941], Right lane) cDNA for 48 hrs and lysed. Equivalent amounts of cell lysates (5 ug per lane) were separated by SDS-PAGE and immunoblotted with anti-SDR9C7. Positive lysates [LY407754] (100ug) and [LC407754] (20ug) can be purchased separately from OriGene.




Immunohistochemical staining of paraffin-embedded Human Kidney tissue within the normal limits using anti-SDR9C7 mouse monoclonal antibody. (Heat-induced epitope retrieval by 10mM citric buffer, pH6.0, 100°C for 10min, TA501723)




Immunohistochemical staining of paraffin-embedded Carcinoma of Human kidney tissue using anti-SDR9C7 mouse monoclonal antibody. (Heat-induced epitope retrieval by 10mM citric buffer, pH6.0, 100°C for 10min, TA501723)



Immunohistochemical staining of paraffin-embedded Human liver tissue within the normal limits using anti-SDR9C7 mouse monoclonal antibody. (Heat-induced epitope retrieval by 10mM citric buffer, pH6.0, 100°C for 10min, TA501723)



Immunohistochemical staining of paraffin-embedded Human bladder tissue within the normal limits using anti-SDR9C7 mouse monoclonal antibody. (Heat-induced epitope retrieval by 10mM citric buffer, pH6.0, 100°C for 10min, TA501723)



HEK293T cells transfected with either [RC210941] overexpress plasmid (Red) or empty vector control plasmid (Blue) were immunostained by anti-SDR9C7 antibody (TA501723), and then analyzed by flow cytometry.