

Product datasheet for **TA502247**


Gephyrin (GPHN) Mouse Monoclonal Antibody [Clone ID: OTI3C6]

Product data:


Product Type:	Primary Antibodies
Clone Name:	OTI3C6
Applications:	FC, IHC, WB
Recommend Dilution:	WB 1:1000~2000, IHC 1:150, FLOW 1:100
Reactivity:	Human
Host:	Mouse
Isotype:	IgG2a
Clonality:	Monoclonal
Immunogen:	Full length human recombinant protein of human GPHN (NP_065857) produced in HEK293T cell.
Formulation:	PBS (PH 7.3) containing 1% BSA, 50% glycerol and 0.02% sodium azide.
Concentration:	0.83 mg/ml
Purification:	Purified from mouse ascites fluids or tissue culture supernatant by affinity chromatography (protein A/G)
Predicted Protein Size:	83.3 kDa
Gene Name:	gephyrin
Database Link:	NP_065857 Entrez Gene 10243 Human
Background:	This gene encodes a neuronal assembly protein that anchors inhibitory neurotransmitter receptors to the postsynaptic cytoskeleton via high affinity binding to a receptor subunit domain and tubulin dimers. In nonneuronal tissues, the encoded protein is also required for molybdenum cofactor biosynthesis. Mutations in this gene may be associated with the neurological condition hyperplexia and also lead to molybdenum cofactor deficiency. Numerous alternatively spliced transcript variants encoding different isoforms have been described; however, the full-length nature of all transcript variants is not currently known. [provided by RefSeq]
Synonyms:	GEPH; GPH; GPHRYN; HKPX1; MOCODC
Protein Families:	Druggable Genome




[View online »](#)

Product images:


HEK293T cells transfected with either [RC205986] overexpress plasmid (Red) or empty vector control plasmid (Blue) were immunostained by anti-GPHN antibody (TA502247), and then analyzed by flow cytometry.




Flow cytometric Analysis of HeLa cells, using anti-GPHN antibody (TA502247), (Red), compared to a nonspecific negative control antibody ([TA50011]), (Blue).




Immunohistochemical staining of paraffin-embedded Human lymphoma tissue using anti-GPHN mouse monoclonal antibody. (Heat-induced epitope retrieval by 10mM citric buffer, pH6.0, 100°C for 10min, TA502247)




Immunohistochemical staining of paraffin-embedded Human lymph node tissue within the normal limits using anti-GPHN mouse monoclonal antibody. (Heat-induced epitope retrieval by 10mM citric buffer, pH6.0, 100°C for 10min, TA502247)




Immunohistochemical staining of paraffin-embedded Carcinoma of Human prostate tissue using anti-GPHN mouse monoclonal antibody. (Heat-induced epitope retrieval by 10mM citric buffer, pH6.0, 100°C for 10min, TA502247)




Immunohistochemical staining of paraffin-embedded Human prostate tissue within the normal limits using anti-GPHN mouse monoclonal antibody. (Heat-induced epitope retrieval by 10mM citric buffer, pH6.0, 100°C for 10min, TA502247)




Immunohistochemical staining of paraffin-embedded Human Kidney tissue within the normal limits using anti-GPHN mouse monoclonal antibody. (Heat-induced epitope retrieval by 10mM citric buffer, pH6.0, 100°C for 10min, TA502247)



Immunohistochemical staining of paraffin-embedded Adenocarcinoma of Human colon tissue using anti-GPHN mouse monoclonal antibody. (Heat-induced epitope retrieval by 10mM citric buffer, pH6.0, 100°C for 10min, TA502247)



HEK293T cells were transfected with the pCMV6-ENTRY control (Left lane) or pCMV6-ENTRY GPHN ([RC205986], Right lane) cDNA for 48 hrs and lysed. Equivalent amounts of cell lysates (5 ug per lane) were separated by SDS-PAGE and immunoblotted with anti-GPHN. Positive lysates [LY402806] (100ug) and [LC402806] (20ug) can be purchased separately from OriGene.



Western blot analysis of extracts (35ug) from 9 different cell lines by using anti-GPHN monoclonal antibody.