

## Product datasheet for TA502923

### GRAP2 Mouse Monoclonal Antibody [Clone ID: OT11G2]


#### Product data:

Product Type:	Primary Antibodies
Clone Name:	OT11G2
Applications:	FC, IHC, WB
Recommend Dilution:	WB 1:500~2000, IHC 1:150, FLOW 1:100
Reactivity:	Human
Host:	Mouse
Isotype:	IgG1
Clonality:	Monoclonal
Immunogen:	Full length human recombinant protein of human GRAP2(NP_004801) produced in HEK293T cell.
Formulation:	PBS (PH 7.3) containing 1% BSA, 50% glycerol and 0.02% sodium azide.
Concentration:	1 mg/ml
Purification:	Purified from mouse ascites fluids or tissue culture supernatant by affinity chromatography (protein A/G)
Predicted Protein Size:	37.7 kDa
Gene Name:	GRB2 related adaptor protein 2
Database Link:	<a href="#">NP_004801 Entrez Gene 9402 Human</a>
Background:	This gene encodes a member of the GRB2/Sem5/Drk family. This member is an adaptor-like protein involved in leukocyte-specific protein-tyrosine kinase signaling. Like its related family member, GRB2-related adaptor protein (GRAP), this protein contains an SH2 domain flanked by two SH3 domains. This protein interacts with other proteins, such as GRB2-associated binding protein 1 (GAB1) and the SLP-76 leukocyte protein (LCP2), through its SH3 domains. Transcript variants utilizing alternative polyA sites exist. [provided by RefSeq]
Synonyms:	GADS; GRAP-2; GRB2L; GRBLG; GrbX; Grf40; GRID; GRPL; Mona; P38
Protein Families:	Druggable Genome
Protein Pathways:	T cell receptor signaling pathway




[View online »](#)


Product images:




HEK293T cells were transfected with the pCMV6-ENTRY control (Left lane) or pCMV6-ENTRY GRAP2 ([RC206546], Right lane) cDNA for 48 hrs and lysed. Equivalent amounts of cell lysates (5 ug per lane) were separated by SDS-PAGE and immunoblotted with anti-GRAP2. Positive lysates [LY401511] (100ug) and [LC401511] (20ug) can be purchased separately from OriGene.




Western blot analysis of extracts (35ug) from 9 different cell lines by using anti-GRAP2 monoclonal antibody.




Immunohistochemical staining of paraffin-embedded Human endometrium tissue within the normal limits using anti-GRAP2 mouse monoclonal antibody. (Heat-induced epitope retrieval by 10mM citric buffer, pH6.0, 100°C for 10min, TA502923)




Immunohistochemical staining of paraffin-embedded Adenocarcinoma of Human endometrium tissue using anti-GRAP2 mouse monoclonal antibody. (Heat-induced epitope retrieval by 10mM citric buffer, pH6.0, 100°C for 10min, TA502923)



Immunohistochemical staining of paraffin-embedded Carcinoma of Human bladder tissue using anti-GRAP2 mouse monoclonal antibody. (Heat-induced epitope retrieval by 10mM citric buffer, pH6.0, 100°C for 10min, TA502923)



HEK293T cells transfected with either [RC206546] overexpress plasmid (Red) or empty vector control plasmid (Blue) were immunostained by anti-GRAP2 antibody (TA502923), and then analyzed by flow cytometry.



Flow cytometric Analysis of Jurkat cells, using anti-GRAP2 antibody (TA502923), (Red), compared to a nonspecific negative control antibody, (Blue).