

## Product datasheet for TA503031

### PDXK Mouse Monoclonal Antibody [Clone ID: OTI3G2]


#### Product data:

Product Type:	Primary Antibodies
Clone Name:	OTI3G2
Applications:	FC, IHC, WB
Recommend Dilution:	WB 1:2000, IHC 1:150, FLOW 1:100
Reactivity:	Human
Host:	Mouse
Isotype:	IgG1
Clonality:	Monoclonal
Immunogen:	Full length human recombinant protein of human PDXK (NP_003672) produced in HEK293T cell.
Formulation:	PBS (PH 7.3) containing 1% BSA, 50% glycerol and 0.02% sodium azide.
Concentration:	0.45 mg/ml
Purification:	Purified from mouse ascites fluids or tissue culture supernatant by affinity chromatography (protein A/G)
Predicted Protein Size:	34.9 kDa
Gene Name:	pyridoxal kinase
Database Link:	<a href="#">NP_003672</a> <a href="#">Entrez Gene 8566 Human</a>
Background:	The protein encoded by this gene phosphorylates vitamin B6, a step required for the conversion of vitamin B6 to pyridoxal-5-phosphate, an important cofactor in intermediary metabolism. The encoded protein is cytoplasmic and probably acts as a homodimer. Alternatively spliced transcript variants have been described, but their biological validity has not been determined. [provided by RefSeq]
Synonyms:	C21orf97; C21orf124; HEL-S-1a; PKH; PNK; PRED79
Protein Families:	Druggable Genome
Protein Pathways:	Metabolic pathways, Vitamin B6 metabolism




[View online »](#)


**Product images:**




HEK293T cells were transfected with the pCMV6-ENTRY control (Left lane) or pCMV6-ENTRY PDXK ([RC200975], Right lane) cDNA for 48 hrs and lysed. Equivalent amounts of cell lysates (5 ug per lane) were separated by SDS-PAGE and immunoblotted with anti-PDXK. Positive lysates [LY418499] (100ug) and [LC418499] (20ug) can be purchased separately from OriGene.




Equivalent amounts of cell lysates (10 ug per lane) of wild-type HEK293T cells (WT, Cat# LC810293T) and PDXK-Knockout HEK293T cells (KO, Cat# [LC841087]) were separated by SDS-PAGE and immunoblotted with anti-PDXK monoclonal antibody TA503031 (1:500). Then the blotted membrane was stripped and reprobed with anti-HSP90 antibody as a loading control.




Immunohistochemical staining of paraffin-embedded Adenocarcinoma of Human breast tissue using anti-PDXK mouse monoclonal antibody. (Heat-induced epitope retrieval by 10mM citric buffer, pH6.0, 100°C for 10min, TA503031)



Immunohistochemical staining of paraffin-embedded Adenocarcinoma of Human ovary tissue using anti-PDXK mouse monoclonal antibody. (Heat-induced epitope retrieval by 10mM citric buffer, pH6.0, 100°C for 10min, TA503031)



Immunohistochemical staining of paraffin-embedded Human pancreas tissue within the normal limits using anti-PDXK mouse monoclonal antibody. (Heat-induced epitope retrieval by 10mM citric buffer, pH6.0, 100°C for 10min, TA503031)



HEK293T cells transfected with either [RC200975] overexpress plasmid (Red) or empty vector control plasmid (Blue) were immunostained by anti-PDXK antibody (TA503031), and then analyzed by flow cytometry.