

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

Product datasheet for TA504171

IGJ (JCHAIN) Mouse Monoclonal Antibody [Clone ID: OTI2B1]

Product data:

Product Type: Primary Antibodies

Clone Name: OTI2B1

Applications: FC, IHC, WB

Recommend Dilution: WB 1:2000, IHC 1:150, FLOW 1:100

Reactivity: Human
Host: Mouse
Isotype: IgG1

Clonality: Monoclonal

Immunogen: Full length human recombinant protein of human IGJ(NP_653247) produced in HEK293T cell.

Formulation: PBS (PH 7.3) containing 1% BSA, 50% glycerol and 0.02% sodium azide.

Concentration: 0.5 mg/ml

Purification: Purified from mouse ascites fluids or tissue culture supernatant by affinity chromatography

(protein A/G)

Predicted Protein Size: 17.9 kDa

Gene Name: joining chain of multimeric IgA and IgM


Database Link: NP 653247 Entrez Gene 3512 Human

Synonyms: IGC|; IG|; |CH






Product images:




HEK293T cells were transfected with the pCMV6-ENTRY control (Left lane) or pCMV6-ENTRY IGJ ([RC207932], Right lane) cDNA for 48 hrs and lysed. Equivalent amounts of cell lysates (5 ug per lane) were separated by SDS-PAGE and immunoblotted with anti-IGJ. Positive lysates [LY403400] (100ug) and [LC403400] (20ug) can be purchased separately from OriGene.




Immunohistochemical staining of paraffinembedded Human lymphoma tissue using anti-IGJ mouse monoclonal antibody. (Heat-induced epitope retrieval by 10mM citric buffer, pH6.0, 100°C for 10min, TA504171)





Immunohistochemical staining of paraffinembedded Human tonsil within the normal limits using anti-IGJ mouse monoclonal antibody. (Heatinduced epitope retrieval by 10mM citric buffer, pH6.0, 100°C for 10min, TA504171)



Flow cytometric Analysis of Jurkat cells, using anti-IGJ antibody (TA504171), (Red), compared to a nonspecific negative control antibody, (Blue).