

Product datasheet for TA504280

Calpain 2 (CAPN2) Mouse Monoclonal Antibody [Clone ID: OTI3G1]


Product data:

Product Type:	Primary Antibodies
Clone Name:	OTI3G1
Applications:	IHC, WB
Recommend Dilution:	WB 1:2000, IHC 1:150
Reactivity:	Human, Dog
Host:	Mouse
Isotype:	IgG2a
Clonality:	Monoclonal
Immunogen:	Full length human recombinant protein of human CAPN2(NP_001739) produced in HEK293T cell.
Formulation:	PBS (PH 7.3) containing 1% BSA, 50% glycerol and 0.02% sodium azide.
Concentration:	1 mg/ml
Purification:	Purified from mouse ascites fluids or tissue culture supernatant by affinity chromatography (protein A/G)
Predicted Protein Size:	79.8 kDa
Gene Name:	calpain 2
Database Link:	NP_001739 Entrez Gene 480118 DogEntrez Gene 824 Human
Background:	The calpains, calcium-activated neutral proteases, are nonlysosomal, intracellular cysteine proteases. The mammalian calpains include ubiquitous, stomach-specific, and muscle-specific proteins. The ubiquitous enzymes consist of heterodimers with distinct large, catalytic subunits associated with a common small, regulatory subunit. This gene encodes the large subunit of the ubiquitous enzyme, calpain 2. Multiple heterogeneous transcriptional start sites in the 5' UTR have been reported. Two transcript variants encoding different isoforms have been found for this gene. [provided by RefSeq, Mar 2009]
Synonyms:	CANP2; CANPL2; CANPml; mCANP
Protein Families:	Druggable Genome, Protease
Protein Pathways:	Alzheimer's disease, Apoptosis, Focal adhesion




[View online »](#)


Product images:




Western blot analysis of extracts (35ug) from 9 different cell lines by using anti-CAPN2 monoclonal antibody (HepG2: human; HeLa: human; SVT2: mouse; A549: human; COS7: monkey; Jurkat: human; MDCK: canine; PC12: rat; MCF7: human) (1:500).




HEK293T cells were transfected with the pCMV6-ENTRY control (Left lane) or pCMV6-ENTRY CAPN2 ([RC205642], Right lane) cDNA for 48 hrs and lysed. Equivalent amounts of cell lysates (5 ug per lane) were separated by SDS-PAGE and immunoblotted with anti-CAPN2. Positive lysates [LY400662] (100ug) and [LC400662] (20ug) can be purchased separately from OriGene.




Immunohistochemical staining of paraffin-embedded Carcinoma of Human pancreas tissue using anti-CAPN2 mouse monoclonal antibody. (Heat-induced epitope retrieval by 10mM citric buffer, pH6.0, 120°C for 3min, TA504280)




Immunohistochemical staining of paraffin-embedded Carcinoma of Human liver tissue using anti-CAPN2 mouse monoclonal antibody. (Heat-induced epitope retrieval by 10mM citric buffer, pH6.0, 120°C for 3min, TA504280)



Immunohistochemical staining of paraffin-embedded Adenocarcinoma of Human colon tissue using anti-CAPN2 mouse monoclonal antibody. (Heat-induced epitope retrieval by 10mM citric buffer, pH6.0, 120°C for 3min, TA504280)



Immunohistochemical staining of paraffin-embedded Adenocarcinoma of Human breast tissue using anti-CAPN2 mouse monoclonal antibody. (Heat-induced epitope retrieval by 10mM citric buffer, pH6.0, 120°C for 3min, TA504280)



Equivalent amounts of cell lysates (10 μ g per lane) of wild-type 293T cells (WT, Cat# LC810293T) and CAPN2-Knockout 293T cells (KO, Cat# [LC841490]) were separated by SDS-PAGE and immunoblotted with anti-CAPN2 monoclonal antibody TA504280, (1:500). Then the blotted membrane was stripped and reprobed with anti- β -actin antibody ([TA811000]) as a loading control.