

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

Product datasheet for TA504524

Carboxypeptidase A (CPA1) Mouse Monoclonal Antibody [Clone ID: OTI1C12]

Product data:

Product Type: Primary Antibodies

Clone Name: OTI1C12
Applications: FC, IF, WB

Recommend Dilution: WB 1:2000, IF 1:100, FLOW 1:100

Reactivity: Human
Host: Mouse
Isotype: IgG1

Clonality: Monoclonal

Immunogen: Full length human recombinant protein of human CPA1(NP_001859) produced in HEK293T

cell

Formulation: PBS (PH 7.3) containing 1% BSA, 50% glycerol and 0.02% sodium azide.

Concentration: 0.82 mg/ml

Purification: Purified from mouse ascites fluids or tissue culture supernatant by affinity chromatography

(protein A/G)

Predicted Protein Size: 45.5 kDa

Gene Name: carboxypeptidase A1

Database Link: NP 001859 Entrez Gene 1357 Human

Background: Three different forms of human pancreatic procarboxypeptidase A have been isolated. This

gene encodes a monomeric pancreatic exopeptidase involved in zymogen inhibition.

[provided by RefSeq]. COMPLETENESS: complete on the 3' end.


Synonyms: CPA

Protein Families: Druggable Genome, ES Cell Differentiation/IPS, Protease, Secreted Protein







Product images:



HEK293T cells were transfected with the pCMV6-ENTRY control (Left lane) or pCMV6-ENTRY CPA1 ([RC202720], Right lane) cDNA for 48 hrs and lysed. Equivalent amounts of cell lysates (5 ug per lane) were separated by SDS-PAGE and immunoblotted with anti-CPA1. Positive lysates [LY419696] (100ug) and [LC419696] (20ug) can be purchased separately from OriGene.



Anti-CPA1 mouse monoclonal antibody (TA504524) immunofluorescent staining of COS7 cells transiently transfected by pCMV6-ENTRY CPA1 ([RC202720]).



HEK293T cells transfected with either [RC202720] overexpress plasmid (Red) or empty vector control plasmid (Blue) were immunostained by anti-CPA1 antibody (TA504524), and then analyzed by flow cytometry.