

Product datasheet for **TA505808**

MAPK4 Mouse Monoclonal Antibody [Clone ID: OTI2D7]


Product data:

Product Type:	Primary Antibodies
Clone Name:	OTI2D7
Applications:	IF, IHC, WB
Recommend Dilution:	WB 1:4000, IHC 1:150, IF 1:100
Reactivity:	Human
Host:	Mouse
Isotype:	IgG2b
Clonality:	Monoclonal
Immunogen:	Full length human recombinant protein of human MAPK4(NP_002738) produced in HEK293T cell.
Formulation:	PBS (PH 7.3) containing 1% BSA, 50% glycerol and 0.02% sodium azide.
Concentration:	1 mg/ml
Purification:	Purified from mouse ascites fluids or tissue culture supernatant by affinity chromatography (protein A/G)
Predicted Protein Size:	65.7 kDa
Gene Name:	mitogen-activated protein kinase 4
Database Link:	NP_002738 Entrez Gene 5596 Human
Synonyms:	ERK-4; ERK4; p63-MAPK; p63MAPK; PRKM4
Protein Families:	Druggable Genome, Protein Kinase




[View online »](#)


Product images:




HEK293T cells were transfected with the pCMV6-ENTRY control (Left lane) or pCMV6-ENTRY MAPK4 ([RC207414], Right lane) cDNA for 48 hrs and lysed. Equivalent amounts of cell lysates (5 ug per lane) were separated by SDS-PAGE and immunoblotted with anti-MAPK4. Positive lysates [LY419131] (100ug) and [LC419131] (20ug) can be purchased separately from OriGene.




Western blot analysis of extracts (10ug) from 8 Human tissue by using anti-MAPK4 monoclonal antibody at 1:200 (1: Testis; 2: Uterus; 3: Breast; 4: Brain; 5: Liver; 6: Ovary; 7: Thyroid gland; 8: Colon).




Immunohistochemical staining of paraffin-embedded Human Kidney tissue within the normal limits using anti-MAPK4 mouse monoclonal antibody. (Heat-induced epitope retrieval by 10mM citric buffer, pH6.0, 120°C for 3min, TA505808)




Immunohistochemical staining of paraffin-embedded Carcinoma of Human kidney tissue using anti-MAPK4 mouse monoclonal antibody. (Heat-induced epitope retrieval by 10mM citric buffer, pH6.0, 120°C for 3min, TA505808)




Immunohistochemical staining of paraffin-embedded Adenocarcinoma of Human ovary tissue using anti-MAPK4 mouse monoclonal antibody. (Heat-induced epitope retrieval by 10mM citric buffer, pH6.0, 120°C for 3min, TA505808)




Immunohistochemical staining of paraffin-embedded Human pancreas tissue within the normal limits using anti-MAPK4 mouse monoclonal antibody. (Heat-induced epitope retrieval by 10mM citric buffer, pH6.0, 120°C for 3min, TA505808)



Immunohistochemical staining of paraffin-embedded Adenocarcinoma of Human endometrium tissue using anti-MAPK4 mouse monoclonal antibody. (Heat-induced epitope retrieval by 10mM citric buffer, pH6.0, 120°C for 3min, TA505808)



Immunohistochemical staining of paraffin-embedded Human tonsil within the normal limits using anti-MAPK4 mouse monoclonal antibody. (Heat-induced epitope retrieval by 10mM citric buffer, pH6.0, 120°C for 3min, TA505808)



Anti-MAPK4 mouse monoclonal antibody (TA505808) immunofluorescent staining of COS7 cells transiently transfected by pCMV6-ENTRY MAPK4 ([RC207414]).