

Product datasheet for **TA506011**

PIK3C2B Mouse Monoclonal Antibody [Clone ID: OTI2G3]

Product data:


Product Type:	Primary Antibodies
Clone Name:	OTI2G3
Applications:	IF, IHC, WB
Recommend Dilution:	WB 1:2000~4000, IHC 1:150, IF 1:100
Reactivity:	Human
Host:	Mouse
Isotype:	IgG2a
Clonality:	Monoclonal
Immunogen:	Full length human recombinant protein of human PIK3C2B(NP_002637) produced in HEK293T cell.
Formulation:	PBS (PH 7.3) containing 1% BSA, 50% glycerol and 0.02% sodium azide.
Concentration:	1 mg/ml
Purification:	Purified from mouse ascites fluids or tissue culture supernatant by affinity chromatography (protein A/G)
Gene Name:	phosphatidylinositol-4-phosphate 3-kinase catalytic subunit type 2 beta
Database Link:	NP_002637 Entrez Gene 5287 Human
Background:	The protein encoded by this gene belongs to the phosphoinositide 3-kinase (PI3K) family. PI3-kinases play roles in signaling pathways involved in cell proliferation, oncogenic transformation, cell survival, cell migration, and intracellular protein trafficking. This protein contains a lipid kinase catalytic domain as well as a C-terminal C2 domain, a characteristic of class II PI3-kinases. C2 domains act as calcium-dependent phospholipid binding motifs that mediate translocation of proteins to membranes, and may also mediate protein-protein interactions. The PI3-kinase activity of this protein is sensitive to low nanomolar levels of the inhibitor wortmanin. The C2 domain of this protein was shown to bind phospholipids but not Ca ²⁺ , which suggests that this enzyme may function in a calcium-independent manner. [provided by RefSeq, Jul 2008]
Synonyms:	C2-PI3K
Protein Families:	Druggable Genome




[View online »](#)

Protein Pathways: Inositol phosphate metabolism, Metabolic pathways, Phosphatidylinositol signaling system


Product images:




HEK293T cells were transfected with the pCMV6-ENTRY control (Left lane) or pCMV6-ENTRY PIK3C2B ([RC218354], Right lane) cDNA for 48 hrs and lysed. Equivalent amounts of cell lysates (5 ug per lane) were separated by SDS-PAGE and immunoblotted with anti-PIK3C2B. Positive lysates [LY419185] (100ug) and [LC419185] (20ug) can be purchased separately from OriGene.



Immunohistochemical staining of paraffin-embedded Adenocarcinoma of Human breast tissue using anti-PIK3C2B mouse monoclonal antibody. (Heat-induced epitope retrieval by 10mM citric buffer, pH6.0, 120°C for 3min, TA506011)



Immunohistochemical staining of paraffin-embedded Human bladder tissue within the normal limits using anti-PIK3C2B mouse monoclonal antibody. (Heat-induced epitope retrieval by 10mM citric buffer, pH6.0, 120°C for 3min, TA506011)



Anti-PIK3C2B mouse monoclonal antibody (TA506011) immunofluorescent staining of COS7 cells transiently transfected by pCMV6-ENTRY PIK3C2B ([RC218354]).