

#### OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

# **Product datasheet for TA807723**

### C Reactive Protein (CRP) Mouse Monoclonal Antibody [Clone ID: OTI4G8]

#### **Product data:**

**Product Type:** Primary Antibodies

Clone Name: OTI4G8
Applications: IHC, WB

**Recommend Dilution:** WB 1:500, IHC 1:150

Reactivity: Human
Host: Mouse
Isotype: IgG1

Clonality: Monoclonal

Immunogen: Full length human recombinant protein of human CRP (NP\_000558) produced in HEK293T

cell.

**Formulation:** PBS (PH 7.3) containing 1% BSA, 50% glycerol and 0.02% sodium azide.

Concentration: 1 mg/ml

**Purification:** Purified from mouse ascites fluids or tissue culture supernatant by affinity chromatography

(protein A/G)

Predicted Protein Size: 23 kDa

Gene Name: C-reactive protein, pentraxin-related

Database Link: NP 000558 Entrez Gene 1401 Human

**Background:** The protein encoded by this gene belongs to the pentaxin family. It is involved in several host

defense related functions based on its ability to recognize foreign pathogens and damaged cells of the host and to initiate their elimination by interacting with humoral and cellular effector systems in the blood. Consequently, the level of this protein in plasma increases greatly during acute phase response to tissue injury, infection, or other inflammatory stimuli.

[provided by RefSeq, Sep 2009]


Synonyms: PTX1

**Protein Families:** Secreted Protein






## **Product images:**



HEK293T cells were transfected with the pCMV6-ENTRY control (Left lane) or pCMV6-ENTRY CRP ([RC217269], Right lane) cDNA for 48 hrs and lysed. Equivalent amounts of cell lysates (5 ug per lane) were separated by SDS-PAGE and immunoblotted with anti-CRP (1:500).



Immunohistochemical staining of paraffinembedded Human tonsil within the normal limits using anti-CRP mouse monoclonal antibody. (Heat-induced epitope retrieval by 1mM EDTA in 10mM Tris buffer (pH8.5) at 120°C for 3min, TA807723) (1:150)