

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

Product datasheet for TA807753

SMCP Mouse Monoclonal Antibody [Clone ID: OTI6G9]

Product data:

Product Type: Primary Antibodies

Clone Name: OTI6G9
Applications: IHC, WB

Recommend Dilution: WB 1:2000, IHC 1:150

Reactivity: Human
Host: Mouse
Isotype: IgG1

Clonality: Monoclonal

Immunogen: Full length human recombinant protein of human SMCP(NP_109588) produced in E.coli.

Formulation: PBS (PH 7.3) containing 1% BSA, 50% glycerol and 0.02% sodium azide.

Concentration: 1 mg/ml

Purification: Purified from mouse ascites fluids or tissue culture supernatant by affinity chromatography

(protein A/G)

Predicted Protein Size: 12.6 kDa

Gene Name: sperm mitochondria associated cysteine rich protein

Database Link: NP 109588 Entrez Gene 4184 Human

Background: Sperm mitochondria differ in morphology and subcellular localization from those of somatic

cells. They are elongated, flattened, and arranged circumferentially to form a helical coiled sheath in the midpiece of the sperm flagellum. The protein encoded by this gene localizes to the capsule associated with the mitochondrial outer membranes and is thought to function in

the organization and stabilization of the helical structure of the sperm's mitochondrial


sheath. [provided by RefSeq, Jul 2008]

Synonyms: HSMCSGEN1; MCS; MCSP






Product images:



HEK293T cells were transfected with the pCMV6-ENTRY control (Left lane) or pCMV6-ENTRY SMCP ([RC205294], Right lane) cDNA for 48 hrs and lysed. Equivalent amounts of cell lysates (5 ug per lane) were separated by SDS-PAGE and immunoblotted with anti-SMCP (1:2000). Positive lysates [LY410741] (100ug) and [LC410741] (20ug) can be purchased separately from OriGene.



Immunohistochemical staining of paraffinembedded Carcinoma of Human bladder tissue using anti-SMCP mouse monoclonal antibody. (Heat-induced epitope retrieval by 1mM EDTA in 10mM Tris buffer (pH8.5) at 120°C for 3min, TA807753) (1:150)