

#### OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

# **Product datasheet for TA808695**

## **CNOT2 Mouse Monoclonal Antibody [Clone ID: OTI6D7]**

#### **Product data:**

**Product Type:** Primary Antibodies

Clone Name: OTI6D7
Applications: IHC

**Recommend Dilution:** IHC 1:150 **Reactivity:** Human

Host: Mouse Isotype: IgG2a

Clonality: Monoclonal

**Immunogen:** Full length human recombinant protein of human CNOT2(NP\_055330) produced in E.coli.

**Formulation:** PBS (PH 7.3) containing 1% BSA, 50% glycerol and 0.02% sodium azide.

Concentration: 1 mg/ml

**Purification:** Purified from mouse ascites fluids or tissue culture supernatant by affinity chromatography

(protein A/G)

**Predicted Protein Size:** 59.6 kDa

**Gene Name:** CCR4-NOT transcription complex subunit 2

Database Link: NP 055330 Entrez Gene 4848 Human

Background: This gene encodes a subunit of the multi-component CCR4-NOT complex. The CCR4-NOT

complex regulates mRNA synthesis and degradation and is also thought to be involved in mRNA splicing, transport and localization. The encoded protein interacts with histone

deacetylases and functions as a repressor of polymerase II transcription. Alternatively spliced

transcript variants have been observed for this gene. [provided by RefSeq, Dec 2010]


Synonyms: CDC36; HSPC131; NOT2; NOT2H

Protein Families: Transcription Factors
Protein Pathways: RNA degradation






## **Product images:**




Immunohistochemical staining of paraffinembedded Adenocarcinoma of Human ovary tissue using anti-CNOT2 mouse monoclonal antibody. (Heat-induced epitope retrieval by 1mM EDTA in 10mM Tris buffer (pH8.5) at 120°C for 3min, TA808695) (1:150)



Immunohistochemical staining of paraffinembedded Carcinoma of Human prostate tissue using anti-CNOT2 mouse monoclonal antibody. (Heat-induced epitope retrieval by 1mM EDTA in 10mM Tris buffer (pH8.5) at 120°C for 3min, TA808695) (1:150)





Immunohistochemical staining of paraffinembedded Carcinoma of Human bladder tissue using anti-CNOT2 mouse monoclonal antibody. (Heat-induced epitope retrieval by 1mM EDTA in 10mM Tris buffer (pH8.5) at 120°C for 3min, TA808695) (1:150)