

## OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

## **Product datasheet for TA808910**

## Myozenin 1 (MYOZ1) Mouse Monoclonal Antibody [Clone ID: OTI4C2]

**Product data:** 

**Product Type:** Primary Antibodies

Clone Name: OTI4C2
Applications: IHC, WB

Recommend Dilution: WB 1:2000, IHC 1:150

Reactivity: Human
Host: Mouse
Isotype: IgG1

Clonality: Monoclonal

**Immunogen:** Human recombinant protein fragment corresponding to amino acids 106-299 of human

MYOZ1(NP\_067068) produced in E.coli.

**Formulation:** PBS (PH 7.3) containing 1% BSA, 50% glycerol and 0.02% sodium azide.

Concentration: 1 mg/ml

**Purification:** Purified from mouse ascites fluids or tissue culture supernatant by affinity chromatography

(protein A/G)

**Predicted Protein Size:** 31.6 kDa **Gene Name:** myozenin 1

Database Link: NP 067068 Entrez Gene 58529 Human

**Background:** The protein encoded by this gene is primarily expressed in the skeletal muscle, and belongs


to the myozenin family. Members of this family function as calcineurin-interacting proteins that help tether calcineurin to the sarcomere of cardiac and skeletal muscle. They play an important role in modulation of calcineurin signaling. [provided by RefSeq, Apr 2012]

**Synonyms:** CS-2; FATZ; MYOZ






## **Product images:**




HEK293T cells were transfected with the pCMV6-ENTRY control (Left lane) or pCMV6-ENTRY MYOZ1 ([RC206448], Right lane) cDNA for 48 hrs and lysed. Equivalent amounts of cell lysates (5 ug per lane) were separated by SDS-PAGE and immunoblotted with anti-MYOZ1 (1:2000). Positive lysates [LY411997] (100ug) and [LC411997] (20ug) can be purchased separately from OriGene.



Immunohistochemical staining of paraffinembedded Adenocarcinoma of Human endometrium tissue using anti-MYOZ1 mouse monoclonal antibody. (Heat-induced epitope retrieval by 1mM EDTA in 10mM Tris buffer (pH8.5) at 120°C for 3min, TA808910) (1:150)





Immunohistochemical staining of paraffinembedded Carcinoma of Human prostate tissue using anti-MYOZ1 mouse monoclonal antibody. (Heat-induced epitope retrieval by 1mM EDTA in 10mM Tris buffer (pH8.5) at 120°C for 3min, TA808910) (1:150)