

#### OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

# **Product datasheet for TA809627**

### OTX2 Mouse Monoclonal Antibody [Clone ID: OTI3G9]

### **Product data:**

**Applications:** 

**Product Type:** Primary Antibodies

WB

Clone Name: OTI3G9

Recommend Dilution: WB 1:2000

Reactivity: Human

Host: Mouse

Isotype: IgG1

Clonality: Monoclonal

**Immunogen:** Full length human recombinant protein of human OTX2 (NP\_068374) produced in E.coli.

**Formulation:** PBS (PH 7.3) containing 1% BSA, 50% glycerol and 0.02% sodium azide.

Concentration: 1 mg/ml

**Purification:** Purified from mouse ascites fluids or tissue culture supernatant by affinity chromatography

(protein A/G)

**Predicted Protein Size:** 32.2 kDa

**Gene Name:** orthodenticle homeobox 2

Database Link: NP 068374 Entrez Gene 5015 Human

**Background:** This gene encodes a member of the bicoid subfamily of homeodomain-containing

transcription factors. The encoded protein acts as a transcription factor and plays a role in brain, craniofacial, and sensory organ development. The encoded protein also influences the proliferation and differentiation of dopaminergic neuronal progenitor cells during mitosis. Mutations in this gene cause syndromic microphthalmia 5 (MCOPS5) and combined pituitary hormone deficiency 6 (CPHD6). This gene is also suspected of having an oncogenic role in medulloblastoma. Alternative splicing results in multiple transcript variants encoding distinct isoforms. Pseudogenes of this gene are known to exist on chromosomes two and nine.

[provided by RefSeq, Jul 2012]

Synonyms: CPHD6; MCOPS5


Protein Families: Embryonic stem cells, Induced pluripotent stem cells, Stem cell - Pluripotency, Transcription

Factors





## **Product images:**



HEK293T cells were transfected with the pCMV6-ENTRY control (Left lane) or pCMV6-ENTRY OTX2 ([RC207479], Right lane) cDNA for 48 hrs and lysed. Equivalent amounts of cell lysates (5 ug per lane) were separated by SDS-PAGE and immunoblotted with anti-OTX2 (1:2000). Positive lysates [LY411946] (100ug) and [LC411946] (20ug) can be purchased separately from OriGene.